

Viscoelastic characteristics of the canine cranial cruciate ligament complex at slow strain rates

Rosti Readioff¹, Brendan Geraghty², Ahmed Elsheikh^{1,3,4} and Eithne Comerford^{2,5}

¹ School of Engineering, University of Liverpool, Liverpool, UK

² Institute of Life Course and Medical Sciences, University of Liverpool, Liverpool, UK

³ Beijing Advanced Innovation Center for Biomedical Engineering, Beihang University, Beijing, China

⁴ UCL Institute of Ophthalmology, NIHR Moorfields BRC, London, UK

⁵ School of Veterinary Science, University of Liverpool, Neston, UK

ABSTRACT

Ligaments including the cruciate ligaments support and transfer loads between bones applied to the knee joint organ. The functions of these ligaments can get compromised due to changes to their viscoelastic material properties. Currently there are discrepancies in the literature on the viscoelastic characteristics of knee ligaments which are thought to be due to tissue variability and different testing protocols. The aim of this study was to characterise the viscoelastic properties of healthy cranial cruciate ligaments (CCLs), from the canine knee (stifle) joint, with a focus on the toe region of the stress-strain properties where any alterations in the extracellular matrix which would affect viscoelastic properties would be seen. Six paired CCLs, from skeletally mature and disease-free Staffordshire bull terrier stifle joints were retrieved as a femur-CCL-tibia complex and mechanically tested under uniaxial cyclic loading up to 10 N at three strain rates, namely 0.1%, 1% and 10%/min, to assess the viscoelastic property of strain rate dependency. The effect of strain history was also investigated by subjecting contralateral CCLs to an ascending (0.1%, 1% and 10%/min) or descending (10%, 1% and 0.1%/min) strain rate protocol. The differences between strain rates were not statistically significant. However, hysteresis and recovery of ligament lengths showed some dependency on strain rate. Only hysteresis was affected by the test protocol and lower strain rates resulted in higher hysteresis and lower recovery. These findings could be explained by the slow process of uncrimping of collagen fibres and the contribution of proteoglycans in the ligament extracellular matrix to intra-fibrillar gliding, which results in more tissue elongations and higher energy dissipation. This study further expands our understanding of canine CCL behaviour, providing data for material models of femur-CCL-tibia complexes, and demonstrating the challenges for engineering complex biomaterials such as knee joint ligaments.

Submitted 18 September 2020

Accepted 2 December 2020

Published 22 December 2020

Corresponding authors

Rosti Readioff, r.readioff@leeds.ac.uk

Eithne Comerford,

eithne.comerford@liverpool.ac.uk

Academic editor

Korakot Nganvongpanit

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.10635

© Copyright

2020 Readioff et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Bioengineering, Veterinary Medicine, Zoology

Keywords Ligaments, Canine knee, Stifle joint, Cruciate ligaments, Viscoelastic, Strain rate, Hysteresis, Recovery

INTRODUCTION

Ligaments play a major role in stifle (knee) joint stability (*Budras, 2007; Levangie & Norkin, 2005*), with part of the primary support being provided by the cranial cruciate ligament (CCL) (*Carpenter & Cooper, 2000; Slatter, 2002*). The CCL is the most commonly ruptured canine stifle joint ligament (CCLR) (*Arnoczky, 1988; Gianotti et al., 2009*) following acute injury or chronic disease, which can lead to destabilisation of surrounding structures and the subsequent development of osteoarthritis (*Arnoczky, 1988; Bennett et al., 1988; Brooks, 2002; Comerford et al., 2006*). There is a large economic cost associated with managing canine CCLR, for example in the United States alone the economic cost was estimated to be at least one billion dollars in 2003 (*Wilke et al., 2005*). Both human and canine CCL failure significantly increases the incidence of age-associated joint degeneration (*Lee et al., 2014; Liu et al., 2003; Peters et al., 2018*) and so understanding the tissue's fundamental material properties can assist with the prediction and effective management of ligament injuries.

The phenomenon of viscoelastic characteristics including strain rate dependency, hysteresis, creep and stress relaxation has been observed consistently in soft biological tissues such as the sclera (*Elsheikh et al., 2010; Geraghty et al., 2020*), cornea (*Elsheikh, Kassem & Jones, 2011; Kazaili, Geraghty & Akhtar, 2019*) and tendon (*Robinson et al., 2004; Zuskov et al., 2020*). Similar to the other biological tissues, ligaments inherit viscoelastic characteristics meaning they exhibiting both elastic and viscous behaviour, hence they are history-and time-dependent (*Bonner et al., 2015; Fung, 1993; Ristianiemi et al., 2018*). The initial part of the non-linear load-deformation behaviour in ligaments is the toe region where the wavy collagen fibres become taut and straighten as load is applied, hence the crimp is removed (*Fratzl et al., 1998*). In this zone, there is a relatively large deformation of the tissue with little increase in load and this permits initial joint deformations with minimal tissue resistance (*Amis, 2004; Dale & Baer, 1974; Fratzl et al., 1998; Wingfield et al., 2000*) (Fig. 1).

Several studies tested for material characteristics of knee joint ligaments at traumatic loading rates and to failure loads (*Crisco, Moore & McGovern, 2002; Crowninshield & Pope, 1976; Lydon et al., 1995*). The loading rate is reported to be directly proportional to the tension which develops in ligaments (*Pioletti, Rakotomanana & Leyvraz, 1999; Woo et al., 1990*). This characteristic was also evident in a study investigating lower strain rates between approximately 2% and 54%/min representing physiological conditions other than trauma (*Haut & Little, 1969*). The study reported no change in the overall shape of the stress-strain curve, however, rapid change in the tangent modulus was found with the slow strain rates (between 1.7% and 10.8 %/min) and the change became progressively smaller with higher strain rates (above 10.8%/min). Similarly, it was reported that strain rate dependency decreases with the increase of deformation rate (*Bonner et al., 2015; Crisco, Moore & McGovern, 2002*). The stress-strain behaviour in the toe region (6% strain) showed strain rate dependency in canine CCL (*Haut & Little, 1969*). However, a study on rabbit medial collateral ligament complexes at varying strain rates (between 0.66% and 9300%/min) showed that the ligaments were only minimally strain rate




Figure 1 A typical stress-strain curve of a knee joint ligament loaded to failure, illustrating the three major regions of the curve. R1: toe region where the ligament fibres tighten and crimp is removed. R2: elastic region. R3: plastic region. The graph is based on previous literature including *Amis (2004)* and *Wingfield et al. (2000)*. Full-size [DOI: 10.7717/peerj.10635/fig-1](https://doi.org/10.7717/peerj.10635/fig-1)

dependent (*Woo et al., 1990; Woo, Gomez & Akeson, 1981*). The small effect of strain rate stiffening could be because the studies combined stress-strain characteristics at the toe region with the elastic region (*Haut & Little, 1969; Ristaniemi et al., 2018*).

During high strain rates, hysteresis (energy dissipated) in the ligament may protect the tissue from injury (*Bonifasi-Lista et al., 2005*). However, there are contradicting findings about hysteresis in soft biological tissues in relation to strain rates. Initially, hysteresis was believed to be weakly dependent on strain rates (*Fung, 1993*). In contrary, a study on the viscoelastic tensile response of bovine cornea showed an increase in hysteresis with decreasing strain rates (*Boyce et al., 2007*). It is suggested that Fung's belief in this phenomenon was based on a small number of experiments on rabbit papillary muscle using only three different strain rates (*Haslach, 2005*). Hence, Fung's findings only approximately support the independence of hysteresis from strain rates.

Therefore, there is a lack of understanding on the strain rate dependency and hysteresis of canine CCLs. Current information is limited to no clear methodological investigations on the strain rate dependency and hysteresis of the CCLs at the toe region (the initial part of non-linear load-deformation behaviour) where collagen fibres tighten and uncrimps with applied load, and importantly, any alterations in the extracellular matrix will be observed (*Comerford et al., 2014; Lujan et al., 2009*). Therefore, the purpose of this study was to characterise the viscoelastic properties of healthy canine CCLs as a femur-CCL-tibia complex, with a focus on the toe region of the stress-strain properties. This quantification is important when comparing the mechanical characteristics of the

CCL and when developing synthetic, auto and allo-grafts to be used in future therapies for ligament replacement.

MATERIALS AND METHODS

CCL storage and preparation

Cadaveric disease-free canine stifle joint pairs from skeletally mature Staffordshire bull terriers ($n = 6$ pairs) euthanatized for reasons other than musculoskeletal injury were obtained with full ethical permission from the Veterinary Research Ethics Committee ((VREC65), Institute of Veterinary Science, University of Liverpool). Inclusion criteria for cadaveric samples were a bodyweight >15 kg and age between 1.5 and 5 years old. The entire stifle joints were frozen at -20°C until required and defrosted at room temperature prior to removing the CCLs as a femur-CCL-tibia complex (Readioff, 2017; Readioff et al., 2020). In order to harvest the femur-CCL-tibia complex, initially the stifle joints were dissected. Subsequently, approximately 10 mm of the femoral and tibial bones were left connected to the CCLs which allowed for the measurement of end-to-end ligament deformation as well as helping to facilitate the clamping of the specimen (Figs. 2 and 3).

The extracted femur-CCL-tibia complexes were maintained in a moistened state in paper towels soaked with phosphate buffered saline (PBS; Sigma, Poole, UK) and frozen at -80°C until they were required for testing (Woo et al., 1986). Prior to testing, the samples were thawed at room temperature and two 1.1 mm arthrodesis wires (Veterinary Instrumentation, Sheffield, UK) were drilled through the tibial and femoral bone ends (Fig. 2). These pins were placed to provide extra grip as well as to replicate the ligament's slight proximal-to-distal outward spiral when secured using custom-built steel clamps (Arnoczky, 1983; Arnoczky & Marshall, 1977). The ducktail clamps were designed to provide a secure grip as well as ensuring that the CCLs were free and unobstructed throughout the experiment (Fig. 3). The clamped samples were then mounted on a mechanical testing machine.

CCL length

A modified version of a previously described method was used to determine the average length of CCL from the craniomedial and caudolateral portions of a ligament (Comerford et al., 2005; Vasseur et al., 1991). In this study, measurements between the insertion and origin of the CCLs at the cranial and caudal planes, as well as the lateral and medial planes were taken using a Vernier callipers (D00352, Duratool, Taiwan, China) accurate to $\pm 10\text{ }\mu\text{m}$. The mean values of these four length measurements were recorded to give an accurate record of the length of the CCL before deformation (Fig. S1).

CCL cross-sectional area

The method by Goodship and Birch was used to measure cross-sectional area (CSA) of the CCLs (Goodship & Birch, 2005). In brief, alginate dental impression paste (UnoDent, UnoDent Ltd., UK) was used to make a mould around the CCL. Once set, the mould was removed from the CCL and was used to create replicas of the CCLs. The replicas were



Figure 2 The extracted cranial cruciate ligaments (CCL) consisted of approximately 10 mm of the femoral and tibial bones forming femur-CCL-tibia complex.

Full-size  DOI: [10.7717/peerj.10635/fig-2](https://doi.org/10.7717/peerj.10635/fig-2)

cut into two in the middle and the surface of the replicas showing middle CSA were estimated using ImageJ (a public domain Java image processing program) (Fig. S2).

Mechanical testing

An Instron 3366 materials testing machine (Instron, Norwood, MA, USA) fitted with a 10 N load cell (Instron 2530-428 with ± 0.025 N accuracy) was used to perform tensile tests. Initially, a preload of 0.1 N was applied to remove laxity within the CCL (Provenzano *et al.*, 2002). Application of the preload was then followed by preconditioning the CCLs to ensure that they were in a steady state and would produce comparable and reproducible load-elongation curves (Butler, Noyes & Grood, 1978; Fung, 1993; Savelberg *et al.*, 1993). Preconditioning involved performing ten loading-unloading cycles up to a maximum load of 10 N at 10 %/min strain rate (Ebrahimi *et al.*, 2019; Woo *et al.*, 1991).




Figure 3 The uniaxial experimental test setup. (A) The design of the custom-made clamps that was used for (B) manufacturing the parts of the experimental testing rig. The custom-built clamps included a (i) cylindrical Perspex tank, (ii) lower and (iii) top grips, (iv) duck-tail sandwich clamps and (v) screws. (C) An example of a cranial cruciate ligament (CCL) clamped and placed in the Perspex tank.

Full-size DOI: [10.7717/peerj.10635/fig-3](https://doi.org/10.7717/peerj.10635/fig-3)

Subsequently the CCL was subjected to cyclic tensile loading-unloading tests investigating stress-strain behaviour of the ligament at the toe region through the application of 10 N load at sequential slow strain rates of 0.1%, 1% and 10%/min. Each strain rate consisted of three loading-unloading cycles which allowed for reproducible results. Between each two cycles, including the preconditioning procedure, a period of 6 min recovery time was given ([Ebrahimi et al., 2019](#); [Viidik, 1968](#)). From the paired stifle joints, the left CCLs were exposed to an ascending strain rate test in which the rate of strain was increased from 0.1% to 1% and to 10%/min and the right CCLs were exposed to a descending strain rate in which the CCL was tested under decreasing strain rates from 10% to 1% and to 0.1 %/min ([Pioletti, Rakotomanana & Leyvraz, 1999](#); [Pioletti & Rakotomanana, 2000](#)). A slow speed was chosen to better observe tissue response to loading at the toe region of load-deformation curves. The reverse orders of strain rate tests (ascending and descending strain rates tests) were carried out to identify characteristics associated with strain history of the ligaments at the toe region.

Data analysis

Analyses on the collected load-deformation data were performed using Microsoft Excel spreadsheets (Microsoft Office 2010, US) and MATLAB (MATLAB R2020a) (code for the analysis and graphs can be found in the [Supplementary Materials](#)). Nominal stress and strain values were estimated following [Eqs. \(1\) and \(2\)](#) ([Haut & Little, 1969](#); [Woo, Gomez & Akeson, 1981](#)) and from these stress and strain values, tangent modulus values were determined ([Eq. 3](#)). Numerical integrations (the trapezoidal rule) on the stress-strain curves were used to estimate the stored energy in the ligaments during loading and unloading tests ([Eq. 4](#)). The hysteresis was then calculated from the difference between the stored energy during loading and unloading tests ([Elsheikh et al., 2008](#)) ([Eq. 5](#)). In addition, ligament extension before and after recovery were studied to investigate strain history and strain rate dependencies as a result of applying loads at different strain rate orders (loading at ascending or descending rates).

$$\sigma = \frac{F}{\text{CSA}} \quad (1)$$

where σ is stress in MPa, F is applied load in N and CSA is cross-sectional area at the middle of the CCL in mm^2 .

$$\varepsilon = \frac{\Delta L}{L_0} \quad (2)$$

where ε is strain, ΔL is change in length in mm ($\Delta L = L_0 - L_1$), L_0 is initial length and L_1 is deformed length of the CCL in mm.

$$E_{\text{tan}} = \frac{\delta \sigma}{\delta \varepsilon} \quad (3)$$

where E_{tan} is tangent modulus in MPa.

$$U = \sum_{k=1}^N \frac{1}{2} \times (\sigma_{k-1} + \sigma_k) \times \Delta \varepsilon_k \quad (4)$$

where U is the stored energy in MPa, N is the resolution of the trapezoidal partition, and $\Delta \varepsilon_k$ is the length of the k th subinterval ($\Delta \varepsilon_k = \varepsilon_k - \varepsilon_{k-1}$).

$$\text{Hysteresis} = U_{\text{Loading}} - U_{\text{Unloading}} \quad (5)$$

where U_{Loading} and $U_{\text{Unloading}}$ represent the stored energy during loading and unloading of the ligaments in MPa.

Statistical analysis

CCL lengths measured at different planes were categorised into cranial, caudal, medial and lateral groups. Statistical tests were performed using one-way analysis of variance (ANOVA) followed by a Bonferroni post-hoc test for multiple comparisons.

A two-tailed t -test (two samples with unequal variance) was used to test for differences between results obtained from the ascending and descending strain rate tests. In addition, one-way ANOVA followed by a Bonferroni post-hoc test for multiple comparisons was performed to test dependencies of tensile responses of the ligaments on strain rates. Distribution of data was illustrated in boxplots and suspected outliers were defined as any value greater than or equal to 1.5 times the interquartile range (range between the first and third quartiles).

All statistical analyses were performed in Microsoft Office Excel and 95% confidence level ($p < 0.05$) was selected to define significance for all statistical tests.

RESULTS

CCL samples

The CCL samples ($n = 6$ paired stifle joints) used to investigate mechanical properties of the ligament were of mixed gender (female = 3 and male = 3) and the bodyweight of the cadavers were in the range of 17–25.5 kg (20.68 ± 3.85 kg).

CCL length

The lengths of the CCLs at different planes were in the range of 7.88–23.16 mm and the measurements at different planes of the individual ligaments are reported in [Table 1](#).

The ANOVA test showed statistically significant results in measuring CCL length in different plane views. The analysis showed that length measurements recorded at different planes were statistically different except for comparisons between medial and lateral planes ([Table S1](#)).

CCL cross-sectional area

The cross-sectional areas of the CCLs were in the range of 11.09–23.62 mm² (16.1 ± 5.1 mm²) and cross-sectional areas of individual ligaments are reported in [Table 2](#).

Table 1 The measured length of cranial cruciate ligaments (CCL) at different measurement planes (cranial, caudal, medial and lateral) for CCLs in paired canine stifle joints ($n = 6$ pairs).

No.	Cranial plane (mm)		Caudal plane (mm)		Medial plane (mm)		Lateral plane (mm)		Average \pm SD (mm)	
	Right	Left	Right	Left	Right	Left	Right	Left	Right	Left
1	13.51	14.54	7.88	8.16	11.76	14.10	11.00	12.31	11.04 \pm 2.35	12.28 \pm 2.91
2	22.79	22.07	11.20	12.00	17.00	13.1	20.54	16.93	17.88 \pm 5.05	16.03 \pm 4.55
3	22.14	22.22	10.21	9.78	17.5	20.36	20.83	19.71	17.67 \pm 5.34	18.02 \pm 5.59
4	21.44	23.16	13.53	11.55	17.51	19.05	16.05	14.37	17.13 \pm 3.31	17.033 \pm 5.12
5	17.88	18.58	10.37	13.02	13.94	15.12	16.51	16.82	14.68 \pm 3.30	15.89 \pm 2.38
6	15.30	17.83	9.20	9.38	13.50	15.81	13.1	12.31	12.78 \pm 2.57	13.83 \pm 3.74
Mean \pm SD (mm)	19.29 \pm 3.47		10.52 \pm 1.79		15.73 \pm 2.60		15.87 \pm 3.34			
Coefficient of variation (%)	18.0		17.0		16.5		21.0			

Table 2 The cross-sectional areas of cranial cruciate ligaments (CCL) for CCLs in paired canine stifle joints ($n = 6$ pairs).

No.	Cross-sectional area (mm ²)	
	Right CCL	Left CCL
1	12.58	14.99
2	14.39	14.41
3	11.09	29.08
4	15.48	13.98
5	12.91	15.69
6	14.93	23.62
Mean \pm SD (mm ²)	16.10 \pm 5.10	
Coefficient of variation (%)	31.7	

Mechanical properties

Stress-strain

The stress-strain curves at 0.1%, 1% and 10%/min strain rates conformed to the typical non-linear behaviour as expected in canine CCLs (Haut & Little, 1969) (Fig. 4A).

The stress-strain curves illustrated an increase in stress with increasing strain, and similarly an increase in stiffness was observed with increasing strain rates (Fig. 4B; Fig. S6). Although there was a small increase in stress with increasing strain rates, the increase was not statistically significant.

The stress responses of the ligaments during ascending test protocol, where the cyclic loading commenced with 0.1%/min then increased to 1%/min and finally to 10%/min, were similar to responses during descending test protocol (the reverse of ascending test protocol). The stress-strain curves show that there are minimal differences in stress values between the two test protocols below 3% strain, and these differences become more distinguishable above 3% strain (Figs. 5A–5C). The testing protocols only minimally affected the stress-strain characteristics and not statistically significant. There are notably




Figure 4 The tensile characteristics of a canine cranial cruciate ligament (CCL) following ascending (Asc) and descending (Desc) protocols at 0.1%, 1% and 10%/min strain rates. (A) A typical cyclic loading and unloading stress-strain curves and (B) tangent modulus-stress behaviour of the loading curves of a CCL at varying strain rates. [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02_img.jpg\) DOI: 10.7717/peerj.10635/fig-4](https://doi.org/10.7717/peerj.10635/fig-4)

different mechanical behaviours among the specimens, as indicated by the grey dots in Figs. 5A–5C, and not all specimens reached 5% strain (Figs. S3–S5).

Tangent modulus-stress

Tangent modulus (E_t), indicating the stiffness behaviour of the CCLs, increased with increasing stress (Fig. 4B) in both ascending and descending testing protocols. Similar to the observations from the stress-strain curves, the tangent modulus-stress lines between the two testing protocols (ascending and descending) were only minimally different and not statistically significant. Although not statistically significant, the increase in tangent modulus with strain rates was notable at higher stress values (Figs. 5D–5F). For example, at 0.5 MPa stress, average tangent modulus values from the ascending test protocol were 26.62, 31.40 and 32.66 MPa during loading at 0.1%, 1% and 10%/min strain rates.

Hysteresis

The results of this study show that hysteresis or dissipated energy are statistically different between the ascending and descending testing protocols ($p = 0.0043$). The mean values for hysteresis at 0.1%, 1% and 10%/min strain rates were 0.0032 (cycle 13), 0.0020 (cycle 16) and 0.0016 (cycle 19) MPa in ascending and 0.0040 (cycle 19), 0.0042 (cycle 16) and 0.0037 (cycle 13) MPa in descending testing protocol (Fig. 6A). The dissipated energy decrease from the first preconditioning cycle to the tenth (last) cycle was 85% for both testing protocols. In addition, hysteresis decreased with increasing strain rates (Fig. 6B). This characteristic was statistically significant during the ascending testing protocol ($p = 0.039$ between 0.1% and 1%/min and $p = 0.013$ between 0.1% and 10%/min). The metadata of hysteresis can be found in (Table S2).




Figure 5 Tensile behaviour of the canine cranial cruciate ligaments was investigated following ascending (red line) and descending (black line) protocols at varying strain rates. The box plots show specimen variation and stress at 1%, 3% and 5% strain during loading at (A) 0.1%/min, (B) 1%/min and (C) 10 %/min strain rates, and tangent modulus at 0.1, 0.3 and 0.5 MPa stress during loading at (D) 0.1%/min, (E) 1%/min and (F) 10%/min strain rates. The outliers are indicated with a red plus sign.

Full-size [DOI: 10.7717/peerj.10635/fig-5](https://doi.org/10.7717/peerj.10635/fig-5)

Recovery

Length of the CCLs before and after the recovery period between each cycle showed consistent values during preconditioning and these values were in the range of




Figure 6 Dissipated energy of the canine cranial cruciate ligaments (CCL) during cyclic loading at varying strain rates. (A) The decrease in mean dissipated energy with increasing cycles of loads. The first ten cycles represent dissipated energy during the precondition stage of the CCLs. From cycles 11–19 dissipated energy values are associated with CCLs during tensile tests at three different strain rates (0.1%, 1% and 10%/min with each test repeated three times). The ascending testing protocol (red line) resulted in a slightly lower dissipated energy compared to the descending (black line) testing protocol. (B) Variations in dissipated energy within the specimens. The median values (indicated by the horizontal line inside the boxes) show a decrease in dissipated energy (hysteresis) with increasing strain rates. However, this observation was only statistically significant (blue line and *) between tensile tests at strain rates of 0.1 and 1%/min, and 0.1 and 10%/min. [Full-size](#) [DOI: 10.7717/peerj.10635/fig-6](#)




Figure 7 Recovery of the canine cranial cruciate ligaments (CCL) during cyclic loading at varying strain rates. (A) The average recovered length of the CCLs at different cycles. During the preconditioning cycles (the first 10 cycles) recovered lengths of the ligaments are similar. Cycles associated with mechanical tests (cycles 11–19) for both testing protocols (ascending in red and descending in black) showed an increase in recovery with increasing strain rates. (B) Variations in length recovery within the CCLs. The box plots show an increase in recovery with increasing strain rates. This characteristic was statistically significant between tensile tests at strain rates of 0.1 and 1%/min, and 0.1 and 10%/min as indicated by the blue asterisk (*) and line. [Full-size](#) [DOI: 10.7717/peerj.10635/fig-7](#)

0.095–0.078 mm during ascending and 0.124–0.095 mm during descending tests (Fig. 7A). Unlike hysteresis, statistical analysis showed that tissue recovery was not different during ascending and descending tests. However, tissue length recovery was strain rate

dependent and statistical analysis showed differences in recovery between 0.1% and 1%/min ($p = 0.018$) and 0.1% and 10%/min ($p = 0.001$) (Fig. 7B).

DISCUSSION

The aim of this study was to gain a greater understanding of the viscoelastic behaviour of the canine CCLs as a femur-CCL-tibia complex at the toe region of the stress-strain curves in order to better mechanically detail the material for its use in developing future therapies. Therefore, we carried out an experimental study investigating the nonlinear viscoelastic properties of CCLs, namely strain rate dependency, hysteresis and recovery, from healthy canine stifle joints. The findings in this study are the first to report the slow strain rate dependency of the canine cruciate ligament across three orders of magnitude with ascending and descending test arrangements. A previous study showed that with high strain rates, the toe region of stress-strain curves appears at lower strain levels (Haut & Little, 1969), however in order to study the toe region in detail without being limited to the level of strain, slow strain rates ($\leq 10\%/min$) were utilised during mechanical tests in the current study.

Similar to previous studies, the measured CCL length and cross-sectional area were used in the calculations for stress and strain values (Wingfield et al., 2000). The difference in the CCLs length in medial and lateral planes was smaller than other planes, and this could be due to the anatomical structure of the femur-CCL-tibia complex.

The non-linear stress-strain pattern for the CCLs is consistent to that previously reported in studies on biological tissues such as tendons and ligaments (Bonner et al., 2015; Crisco, Moore & McGovern, 2002; Haut & Little, 1969; Pioletti, Rakotomanana & Leyvraz, 1999; Pioletti & Rakotomanana, 2000). The stress-strain and tangent modulus-stress characteristics of the CCLs were similar during the ascending and descending testing protocols (Figs. 4 and 5). These findings are similar to a previous study on bovine anterior cruciate ligament-bone complex where specimens were loaded up to 300 N at seven different strain rates (6%, 60%, 300%, 600%, 1,200%, 1,800% and 2,400%/min) and then tested for strain rate order by reloading the ligaments at the 6% and 300%/min strain rates (Pioletti, Rakotomanana & Leyvraz, 1999). They found identical stress-strain behaviour for the initial and reloaded specimens suggesting no difference in changing test protocols via strain rate orders. Their study applied higher strain rates (6–2,400%/min) than those used in the current paper and they reloaded the tissue in an ascending strain rate order only. Pioletti, Rakotomanana & Leyvraz (1999) did not study tissue hysteresis or recovery, but they reported increases in linear tangent moduli with increasing strain rates (although not statistically significant) which is similar to the findings in the current study.

The mechanical response of human knee ligaments to loading depends on strain rate which is less pronounced at lower rates (Dorlot et al., 1980; Van Dommelen et al., 2005) and this was also observed in the current study. It is believed that during lower strain rates the collagen fibrils in patella tendons undergo significantly less recruitment (Clemmer et al., 2010) and this could potentially be similar in the case of the CCLs. At slow strain rates ($\leq 10\%/min$) the collagen fibrils uncrimp with applied load and then show

intra-fibrillar gliding (Bonner *et al.*, 2015; Karunaratne, Li & Bull, 2018). However, at fast strain rates ($\geq 300\%/min$) fibrils go from an unloaded state directly to intra-fibrillar gliding where the matrix bond between the collagen molecules are broken before the removal of collagen crimps (Bonner *et al.*, 2015). This could mean that the extracellular matrix components such as proteoglycans, which is directly linked to the mechanics of ligaments during uncrimping of the collagen fibres, might not affect the mechanical response during loads at higher strain rates.

In our study, hysteresis, which represents the dissipation of energy within the tissue, has shown some dependencies on strain rates and decreases with increasing strain rates (Fig. 6B). This finding contradicts the conclusions from previous literature (Bonifasi-Lista *et al.*, 2005; Woo, Gomez & Akeson, 1981) but agrees with a recent study on tendon fascicle mechanics (Rosario & Roberts, 2020) and a study on bovine cornea (Boyce *et al.*, 2007) which found a decrease in strain energy storage with increased loading rate. The discrepancies in results might be due to following different testing protocols, in particular the rate of applied loads. In the current study, where slow strain rates of $\leq 10\%/min$ were used, the tissue goes through more steps (uncrimping collagen fibres, intra-fibrillar gliding and then loading of collagen respectively with applied loads) during lower strain rates (Bonner *et al.*, 2015), and this slow process results in more tissue elongations hence higher energy dissipation. Energy dissipation was highest during the first precondition cycle and this could be a result of tissue handling and loading history of the CCLs (Fig. 6). In addition, it is possible that the ligament was dissipating higher energy at the initial loading cycles (during preconditioning) because of microstructural reorganisation. Similar to previous literature, this study found a decrease in hysteresis during preconditioning cycles (Woo *et al.*, 1986; Yahia & Drouin, 1990).

The CCLs showed higher length recovery during higher strain rates compared to the lower strain rates (Fig. 7). This behaviour could be a result of resilience in the fascicular level of the tissue. It has been reported that during slow strain rates, the resilience in the fascicular level is lowest (Rosario & Roberts, 2020) which could lead to higher changes in the microstructural organisation of the tissue. Hence, the tissue's length might not fully recover within the same recovery time as during higher strain rate tests. However, it is important to note that although the higher strain rate might seem to result in a more recovered ligament length, it is possible that the tissue collagen fibres are still crimping back as a result of previous loading history or insufficient recovery time. Further study in this area especially at the microstructural level of knee ligaments is necessary to better understand the effects of loading rates on the organisations of the fibres and extracellular matrix.

There were several limitations to our study. Preparing specimens for mechanical tests as a whole unit (femur-CCL-tibia complex) might have introduced some limitations such as overlooking the complexity of the anatomical structure of the CCLs which consists of two fibre bundles (caudolateral (CLB) and craniomedial bands (CMB)) functioning independently from one another in stifle joint flexion and extension (Arnoczky & Marshall, 1977; Carpenter & Cooper, 2000). Independent functioning of the CLB and the CMB allows the fibre bundles to reach their maximum potential (Arnoczky & Marshall,

1977; Carpenter & Cooper, 2000; Tanegashima et al., 2019). However, it is important to note that these two fibre bundles are not structurally segregated within the tissue, thus allowing the ligament to function as a united structure (Heffron & Campbell, 1978). In addition, the approximation methods adopted to measure the cross-sectional area and length of the specimens might be considered as another limitation. Further investigation with a larger number of specimens might improve the reliability of the statistical analysis and provide a broader view on the effect of cadaveric demography (i.e., age, gender, bodyweight) on the mechanical properties and microstructural organisations of knee ligaments (Duval et al., 1999; Woo, Ohland & Weiss, 1990; Woo et al., 1990). In addition, investigating material properties of the ligaments using other quantitative measures such as stress relaxation or creep could further expand our knowledge on the viscoelasticity of the tissue (Amis, 1985).

CONCLUSIONS

The current study focused on the viscoelastic behaviour, such as strain rate dependency, hysteresis and recovery of canine CCLs at slow strain rates to better understand the tissue behaviour at the toe region where the constituents of the extracellular matrix makes a major contribution to ligament mechanics.

Our changing test protocols via strain rate orders only affected hysteresis which might be a result of the strain history of the tissue or high-level of biological variability across samples (Gardiner & Weiss, 2003; Harris et al., 2016). The stress-strain of the CCLs at the toe-region associated with the extracellular matrix of the ligaments was not strain rate dependent. However, hysteresis and recovery were strain rate dependent and this is likely due to changes in microstructural organisation of the ligaments during mechanical tests.

The result of our study indicates the need for further investigations on the viscoelastic behaviour of the canine CCLs when loaded with different orders of strain rates, with a focus on extracellular matrix and collagen fibre organisations.

ACKNOWLEDGEMENTS

We thank Mr. Lee Moore, Mr. Ben Jones and the staff at Veterinary Teaching Suite, School of Veterinary Science for their assistance during sample collection. We also thank Mr. John Curran at School of Engineering, University of Liverpool, for their assistance during manufacturing parts of the experimental setup.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the School of Engineering at the University of Liverpool (Rosti Readioff), the Wellcome Trust Institutional Strategic Support Fund (WT 204822/Z/16/Z) (Eithne Comerford), and by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Moorfields Eye Hospital NHS Foundation Trust

and UCL Institute of Ophthalmology (Ahmed Elsheikh). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

School of Engineering at the University of Liverpool (Rosti Readioff).

Wellcome Trust Institutional Strategic Support Fund: WT 204822/Z/16/Z (Eithne Comerford).

National Institute for Health Research (NIHR) Biomedical Research Centre based at Moorfields Eye Hospital NHS Foundation Trust.

UCL Institute of Ophthalmology (Ahmed Elsheikh).

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Rosti Readioff conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Brendan Geraghty analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Ahmed Elsheikh conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Eithne Comerford conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Cadaveric canine stifle joints were obtained with full ethical permission from the Veterinary Research Ethics Committee ((VREC65), Institute of Veterinary Science, University of Liverpool, Liverpool, UK).

Data Availability

The following information was supplied regarding data availability:

Raw, analysed and statistical data are available at Zenodo:

Readioff, Rosti. (2020). Raw data for viscoelastic anterior cruciate ligament characteristics (Version v2.0) (Data set). Zenodo. DOI [10.5281/zenodo.4264834](https://doi.org/10.5281/zenodo.4264834).

Codes used for the initial data analysis and plotting the graphs are available at Zenodo:

Rosti Readioff. (2020, November 9). RostiReadioff/RReadioff_Viscoelasticity_of_Knee_Ligaments: Codes for plotting and analysing VE ACL data (Version v2.0). Zenodo. DOI [10.5281/zenodo.4264881](https://doi.org/10.5281/zenodo.4264881).

Additional figures and tables are available as a [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.10635#supplemental-information>.

REFERENCES

- Amis AA. 1985. Biomechanics of ligaments. In: *Ligament Injuries and their Treatment*. London: Chapman and Hall Ltd, 3–25.
- Amis AA. 2004. The biomechanics of ligaments. In: Poitout DG, ed. *Biomechanics and Biomaterials in Orthopedics*. London: Springer, 550–563.
- Arnoczky SP. 1983. Anatomy of the anterior cruciate ligament. *Clinical Orthopaedics and Related Research* 172:19–25.
- Arnoczky SP. 1988. The cruciate ligaments: the enigma of the canine stifle. *Journal of Small Animal Practice* 29(2):71–90 DOI 10.1111/j.1748-5827.1988.tb02267.x.
- Arnoczky SP, Marshall JL. 1977. The cruciate ligaments of the canine stifle: An anatomical and functional analysis. *American Journal of Veterinary Research* 38:1807–1814.
- Bennett D, Tennant B, Lewis DG, Baughan J, May C, Carter S. 1988. A reappraisal of anterior cruciate ligament disease in the dog. *Journal of Small Animal Practice* 29(5):275–297 DOI 10.1111/j.1748-5827.1988.tb02286.x.
- Bonifasi-Lista C, Lake SP, Small MS, Weiss JA. 2005. Viscoelastic properties of the human medial collateral ligament under longitudinal, transverse and shear loading. *Journal of Orthopaedic Research* 23(1):67–76 DOI 10.1016/j.orthres.2004.06.002.
- Bonner TJ, Newell N, Karunaratne A, Pullen AD, Amis AA, Bull AMJ, Masouros SD. 2015. Strain-rate sensitivity of the lateral collateral ligament of the knee. *Journal of the Mechanical Behavior of Biomedical Materials* 41:261–270 DOI 10.1016/j.jmbbm.2014.07.004.
- Boyce BL, Jones RE, Nguyen TD, Grazier JM. 2007. Stress-controlled viscoelastic tensile response of bovine cornea. *Journal of Biomechanics* 40(11):2367–2376 DOI 10.1016/j.jbiomech.2006.12.001.
- Brooks PM. 2002. Impact of osteoarthritis on individuals and society: how much disability? Social consequences and health economic implications. *Current Opinion in Rheumatology* 14(5):573–577 DOI 10.1097/00002281-200209000-00017.
- Budras K-D. 2007. *Anatomy of the dog, Vet series*. Fifth Edition. Hannover: Schlütersche.
- Butler DL, Noyes FR, Grood ES. 1978. Measurement of the mechanical properties of ligaments. In: *CRC Handbook of Engineering in Medicine and Biology Section B: Instruments and Measurements*. West Palm Beach: CRC Press Inc., 279–314.
- Carpenter DH, Cooper RC. 2000. Mini review of canine stifle joint anatomy. *Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C* 29(6):321–329 DOI 10.1046/j.1439-0264.2000.00289.x.
- Clemmer J, Liao J, Davis D, Horstemeyer MF, Williams LN. 2010. A mechanistic study for strain rate sensitivity of rabbit patellar tendon. *Journal of Biomechanics* 43(14):2785–2791 DOI 10.1016/j.jbiomech.2010.06.009.
- Comerford EJ, Geraghty B, Hama Rashid R, Elsheikh A. 2014. The contribution of proteoglycans to the viscoelasticity of the canine anterior cruciate ligament. *Osteoarthritis Cartilage* 22:S313.
- Comerford EJ, Tarlton JF, Innes JF, Johnson KA, Amis AA, Bailey AJ. 2005. Metabolism and composition of the canine anterior cruciate ligament relate to differences in knee joint mechanics and predisposition to ligament rupture. *Journal of Orthopaedic Research* 23(1):61–66 DOI 10.1016/j.orthres.2004.05.016.

- Comerford EJ, Tarlton JF, Wales A, Bailey AJ, Innes JF. 2006.** Ultrastructural differences in cranial cruciate ligaments from dogs of two breeds with a differing predisposition to ligament degeneration and rupture. *Journal of Comparative Pathology* **134**(1):8–16 DOI [10.1016/j.jcpa.2005.06.004](https://doi.org/10.1016/j.jcpa.2005.06.004).
- Crisco JJ, Moore DC, McGovern RD. 2002.** Strain-rate sensitivity of the rabbit MCL diminishes at traumatic loading rates. *Journal of Biomechanics* **35**(10):1379–1385 DOI [10.1016/S0021-9290\(02\)00167-7](https://doi.org/10.1016/S0021-9290(02)00167-7).
- Crowninshield RD, Pope MH. 1976.** The strength and failure characteristics of rat medial collateral ligaments. *Journal of Trauma: Injury, Infection, and Critical Care* **16**(2):99–105 DOI [10.1097/00005373-197602000-00004](https://doi.org/10.1097/00005373-197602000-00004).
- Dale WC, Baer E. 1974.** Fiber-buckling in composite systems: a model for ultrastructure of uncalcified collagen tissues. *Journal of Materials Science* **9**(3):369–382 DOI [10.1007/BF00737836](https://doi.org/10.1007/BF00737836).
- Dorlot JM, Ait Ba Sidi M, Tremblay GM, Drouin G. 1980.** Load elongation behavior of the canine anterior cruciate ligament. *Journal of Biomechanical Engineering* **102**(3):190–193 DOI [10.1115/1.3149572](https://doi.org/10.1115/1.3149572).
- Duval JM, Budsberg SC, Flo GL, Sammarco JL. 1999.** Breed, sex, and body weight as risk factors for rupture of the cranial cruciate ligament in young dogs. *Journal of the American Veterinary Medical Association* **215**:811–814.
- Ebrahimi M, Mohammadi A, Ristaniemi A, Stenroth L, Korhonen RK. 2019.** The effect of different preconditioning protocols on repeatability of bovine ACL stress-relaxation response in tension. *Journal of the Mechanical Behavior of Biomedical Materials* **90**:493–501 DOI [10.1016/j.jmbbm.2018.10.041](https://doi.org/10.1016/j.jmbbm.2018.10.041).
- Elsheikh A, Geraghty B, Alhasso D, Knappett J, Campanelli M, Rama P. 2010.** Regional variation in the biomechanical properties of the human sclera. *Experimental Eye Research* **90**(5):233–624 DOI [10.1016/j.exer.2010.02.010](https://doi.org/10.1016/j.exer.2010.02.010).
- Elsheikh A, Kassem W, Jones SW. 2011.** Strain-rate sensitivity of porcine and ovine corneas. *Acta of Bioengineering and Biomechanics* **13**:25–36.
- Elsheikh A, Wang D, Rama P, Campanelli M, Garway-Heath D. 2008.** Experimental assessment of human corneal hysteresis. *Current Eye Research* **33**(3):205–213 DOI [10.1080/02713680701882519](https://doi.org/10.1080/02713680701882519).
- Fratzl P, Misof K, Zizak I, Rapp G, Amenitsch H, Bernstorff S. 1998.** Fibrillar structure and mechanical properties of collagen. *Journal of Structural Biology* **122**(1–2):119–122 DOI [10.1006/jsbi.1998.3966](https://doi.org/10.1006/jsbi.1998.3966).
- Fung YC. 1993.** *Biomechanics: mechanical properties of living tissues*. Second Edition. New York: Springer.
- Gardiner JC, Weiss JA. 2003.** Subject-specific finite element analysis of the human medial collateral ligament during valgus knee loading. *Journal of Orthopaedic Research* **21**(6):1098–1106 DOI [10.1016/S0736-0266\(03\)00113-X](https://doi.org/10.1016/S0736-0266(03)00113-X).
- Geraghty B, Abass A, Eliasy A, Jones SW, Rama P, Kassem W, Akhtar R, Elsheikh A. 2020.** Inflation experiments and inverse finite element modelling of posterior human sclera. *Journal of Biomechanics* **98**:109438 DOI [10.1016/j.jbiomech.2019.109438](https://doi.org/10.1016/j.jbiomech.2019.109438).
- Gianotti SM, Marshall SW, Hume PA, Bunt L. 2009.** Incidence of anterior cruciate ligament injury and other knee ligament injuries: a national population-based study. *Journal of Science and Medicine in Sport* **12**(6):622–627 DOI [10.1016/j.jsams.2008.07.005](https://doi.org/10.1016/j.jsams.2008.07.005).

- Goodship AE, Birch HL. 2005. Cross sectional area measurement of tendon and ligament in vitro: a simple, rapid, non-destructive technique. *Journal of Biomechanics* 38(3):605–608 DOI 10.1016/j.jbiomech.2004.05.003.
- Harris MD, Cyr AJ, Ali AA, Fitzpatrick CK, Rullkoetter PJ, Maletsky LP, Shelburne KB. 2016. A combined experimental and computational approach to subject-specific analysis of knee joint laxity. *Journal of Biomechanical Engineering* 138(8):1019 DOI 10.1115/1.4033882.
- Haslach HW. 2005. Nonlinear viscoelastic, thermodynamically consistent, models for biological soft tissue. *Biomechanics and Modeling in Mechanobiology* 3(3):172–189 DOI 10.1007/s10237-004-0055-6.
- Haut RC, Little RW. 1969. Rheological properties of canine anterior cruciate ligaments. *Journal of Biomechanics* 2(3):289–298 DOI 10.1016/0021-9290(69)90085-2.
- Heffron LE, Campbell JR. 1978. Morphology, histology and functional anatomy of the canine cranial cruciate ligament. *Veterinary Record* 102(13):280–283 DOI 10.1136/vr.102.13.280.
- Karunaratne A, Li S, Bull AMJ. 2018. Nano-scale mechanisms explain the stiffening and strengthening of ligament tissue with increasing strain rate. *Scientific Reports* 8:3707 DOI 10.1038/s41598-018-21786-z.
- Kazaili A, Geraghty B, Akhtar R. 2019. Microscale assessment of corneal viscoelastic properties under physiological pressures. *Journal of the Mechanical Behavior of Biomedical Materials* 100:103375 DOI 10.1016/j.jmbbm.2019.103375.
- Lee K, Williamson K, Clegg P, Comerford E, Canty-Laird E. 2014. The stem cell niche in tendon and ligament: investigating alterations with ageing and disease. *British Journal of Sports Medicine—BMJ* 48(Suppl. 2):A35–A36 DOI 10.1136/bjsports-2014-094114.54.
- Levangie PK, Norkin CC. 2005. *Joint structure and function: a comprehensive analysis*. Fourth Edition. Philadelphia: F.A. Davis Co.
- Liu WH, Burton-Wurster N, Glant TT, Tashman S, Sumner DR, Kamath RV, Lust G, Kimura JH, Cs-Szabo G. 2003. Spontaneous and experimental osteoarthritis in dog: similarities and differences in proteoglycan levels. *Journal of Orthopaedic Research* 21(4):730–737 DOI 10.1016/S0736-0266(03)00002-0.
- Lujan TJ, Underwood CJ, Jacobs NT, Weiss JA. 2009. Contribution of glycosaminoglycans to viscoelastic tensile behavior of human ligament. *Journal of Applied Physiology* 106(2):423–431 DOI 10.1152/jappphysiol.90748.2008.
- Lydon C, Crisco JJ, Panjabi M, Galloway M. 1995. Effect of elongation rate on the failure properties of the rabbit anterior cruciate ligament. *Clinical Biomechanics* 10(8):428–433 DOI 10.1016/0268-0033(95)00019-2.
- Peters AE, Akhtar R, Comerford EJ, Bates KT. 2018. The effect of ageing and osteoarthritis on the mechanical properties of cartilage and bone in the human knee joint. *Scientific Reports* 8(1):5931 DOI 10.1038/s41598-018-24258-6.
- Pioletti DP, Rakotomanana LR. 2000. Non-linear viscoelastic laws for soft biological tissues. *European Journal of Mechanics - A/Solids* 19(5):749–759 DOI 10.1016/S0997-7538(00)00202-3.
- Pioletti DP, Rakotomanana LR, Leyvraz PF. 1999. Strain rate effect on the mechanical behavior of the anterior cruciate ligament–bone complex. *Medical Engineering & Physics* 21(2):95–100 DOI 10.1016/S1350-4533(99)00028-4.
- Provenzano PP, Heisey D, Hayashi K, Lakes R, Vanderby R Jr. 2002. Subfailure damage in ligament: a structural and cellular evaluation. *Journal of Applied Physiology* 92(1):362–371 DOI 10.1152/jappl.2002.92.1.362.
- Radioff R. 2017. *Viscoelastic behaviour of the canine cranial cruciate ligament complex*. Liverpool: University of Liverpool.

- Radioff R, Geraghty B, Comerford E, Elsheikh A. 2020.** A full-field 3D digital image correlation and modelling technique to characterise anterior cruciate ligament mechanics ex vivo. *Acta Biomaterialia* **113**:417–428 DOI [10.1016/j.actbio.2020.07.003](https://doi.org/10.1016/j.actbio.2020.07.003).
- Ristaniemi A, Stenroth L, Mikkonen S, Korhonen RK. 2018.** Comparison of elastic, viscoelastic and failure tensile material properties of knee ligaments and patellar tendon. *Journal of Biomechanics* **79**:31–38 DOI [10.1016/j.jbiomech.2018.07.031](https://doi.org/10.1016/j.jbiomech.2018.07.031).
- Robinson PS, Lin TW, Reynolds PR, Derwin KA, Iozzo RV, Soslowsky LJ. 2004.** Strain-rate sensitive mechanical properties of tendon fascicles from mice with genetically engineered alterations in collagen and decorin. *Journal of Biomechanical Engineering* **126**(2):252–257 DOI [10.1115/1.1695570](https://doi.org/10.1115/1.1695570).
- Rosario MV, Roberts TJ. 2020.** Loading rate has little influence on tendon fascicle mechanics. *Frontiers in Physiology* **11**:1247 DOI [10.3389/fphys.2020.00255](https://doi.org/10.3389/fphys.2020.00255).
- Savelberg HHCM, Kooloos JGM, Huiskes R, Kauer JMG. 1993.** An indirect method to assess wrist ligament forces with particular regard to the effect of preconditioning. *Journal of Biomechanics* **26**(11):1347–1351 DOI [10.1016/0021-9290\(93\)90358-L](https://doi.org/10.1016/0021-9290(93)90358-L).
- Slatyer DH. 2002.** Textbook of small animal surgery. Third Edition. Philadelphia: Saunders.
- Tanegashima K, Edamura K, Akita Y, Yamazaki A, Yasukawa S, Seki M, Asano K, Nakayama T, Katsura T, Hayashi K. 2019.** Functional anatomy of the craniomedial and caudolateral bundles of the cranial cruciate ligament in beagle dogs. *Veterinary and Comparative Orthopaedics and Traumatology* **32**(3):182–191 DOI [10.1055/s-0039-1678711](https://doi.org/10.1055/s-0039-1678711).
- Van Dommelen JAW, Ivarsson BJ, Jolandan MM, Millington SA, Raut M, Kerrigan JR, Crandall JR, Diduch DR. 2005.** Characterization of the rate-dependent mechanical properties and failure of human knee ligaments. *SAE International Journal of Engines* **114**:80–90.
- Vasseur PB, Stevenson S, Gregory CR, Rodrigo JJ, Pauli S, Heitter D, Sharkey N. 1991.** Anterior cruciate ligament allograft transplantation in dogs. *Clinical Orthopaedics and Related Research* **269**:295–304 DOI [10.1097/00003086-199108000-00040](https://doi.org/10.1097/00003086-199108000-00040).
- Viidik A. 1968.** Elasticity and tensile strength of anterior cruciate ligament in rabbits as influenced by training. *Acta Physiologica Scandinavica* **74**(3):372–380 DOI [10.1111/j.1365-201X.1968.tb10930.x](https://doi.org/10.1111/j.1365-201X.1968.tb10930.x).
- Wilke VL, Robinson DA, Evans RB, Rothschild MF, Conzemius MG. 2005.** Estimate of the annual economic impact of treatment of cranial cruciate ligament injury in dogs in the United States. *Journal of the American Veterinary Medical Association* **227**(10):1604–1607 DOI [10.2460/javma.2005.227.1604](https://doi.org/10.2460/javma.2005.227.1604).
- Wingfield C, Amis AA, Stead AC, Law HT. 2000.** Comparison of the biomechanical properties of rottweiler and racing greyhound cranial cruciate ligaments. *Journal of Small Animal Practice* **41**(7):303–307 DOI [10.1111/j.1748-5827.2000.tb03206.x](https://doi.org/10.1111/j.1748-5827.2000.tb03206.x).
- Woo SLY, Gomez MA, Akeson WH. 1981.** The time and history-dependent viscoelastic properties of the canine medial collateral ligament. *Journal of Biomechanical Engineering* **103**(4):293–298 DOI [10.1115/1.3138295](https://doi.org/10.1115/1.3138295).
- Woo SLY, Hollis JM, Adams DJ, Lyon RM, Takai S. 1991.** Tensile properties of the human femur-anterior cruciate ligament-tibia complex: the effects of specimen age and orientation. *American Journal of Sports Medicine* **19**(3):217–225 DOI [10.1177/036354659101900303](https://doi.org/10.1177/036354659101900303).
- Woo SLY, Ohland KJ, Weiss JA. 1990.** Aging and sex-related changes in the biomechanical properties of the rabbit medial collateral ligament. *Mechanisms of Ageing and Development* **56**(2):129–142 DOI [10.1016/0047-6374\(90\)90004-Y](https://doi.org/10.1016/0047-6374(90)90004-Y).

- Woo SLY, Orlando CA, Camp JF, Akeson WH. 1986.** Effects of postmortem storage by freezing on ligament tensile behavior. *Journal of Biomechanics* **19**(5):399–404
[DOI 10.1016/0021-9290\(86\)90016-3](https://doi.org/10.1016/0021-9290(86)90016-3).
- Woo SLY, Peterson RH, Ohland KJ, Sites TJ, Danto MI. 1990.** The effects of strain rate on the properties of the medial collateral ligament in skeletally immature and mature rabbits: a biomechanical and histological study. *Journal of Orthopaedic Research* **8**(5):712–721
[DOI 10.1002/jor.1100080513](https://doi.org/10.1002/jor.1100080513).
- Yahia LH, Drouin G. 1990.** Study of the hysteresis phenomenon in canine anterior cruciate ligaments. *Journal of Biomedical Engineering* **12**(1):57–62
[DOI 10.1016/0141-5425\(90\)90116-5](https://doi.org/10.1016/0141-5425(90)90116-5).
- Zuskov A, Freedman BR, Gordon JA, Sarver JJ, Buckley MR, Soslowsky LJ. 2020.** Tendon biomechanics and crimp properties following fatigue loading are influenced by tendon type and age in mice. *Journal of Orthopaedic Research* **38**(1):36–42 [DOI 10.1002/jor.24407](https://doi.org/10.1002/jor.24407).